



Contents lists available at ScienceDirect

Vaccine

journal homepage: www.elsevier.com/locate/vaccine

Evaluating public acceptability of a potential Lyme disease vaccine using a population-based, cross-sectional survey in high incidence areas of the United States

Sarah A. Hook^{a,*}, AmberJean P. Hansen^b, Sara A. Niesobecki^b, James I. Meek^b, Jenna K.H. Bjork^c, Erin M. Kough^c, Molly S. Peterson^c, Elizabeth K. Schiffman^c, Heather J. Rutz^d, Adam J. Rowe^e, Jennifer L. White^e, Jennifer L. Peel^f, Brad J. Biggerstaff^a, Alison F. Hinckley^a

^a Centers for Disease Control and Prevention, Division of Vector-borne Diseases, 3156 Rampart Rd., Fort Collins, CO 80521, USA

^b Connecticut Emerging Infections Program, Yale School of Public Health, One Church Street, 7th Floor, New Haven, CT 06510, USA

^c Minnesota Department of Health, 625 Robert St N, St Paul, MN 55164, USA

^d Maryland Department of Health, 201 W. Preston Street, 3rd Floor, Baltimore, MD 21201, USA

^e New York State Department of Health, Empire Plaza, Corning Tower, State St, Albany, NY 12203, USA

^f Colorado State University, Environmental Health Building, 350 W. Lake St., Fort Collins, CO 80523, USA

ARTICLE INFO

Article history:

Received 12 August 2021

Received in revised form 9 November 2021

Accepted 22 November 2021

Available online xxx

Keywords:

Tick
Tickborne
Lyme disease
Vaccine
Acceptability

ABSTRACT

Background: Lyme disease incidence is increasing, despite current prevention options. New Lyme disease vaccine candidates are in development, however, investigation of the acceptability of a Lyme disease vaccine among potential consumers is needed prior to any vaccine coming to market. We conducted a population-based, cross-sectional study to estimate willingness to receive a potential Lyme disease vaccine and factors associated with willingness.

Methods: The web-based survey was administered to a random sample of Connecticut, Maryland, Minnesota, and New York residents June–July 2018. Survey-weighted descriptive statistics were conducted to estimate the proportion willing to receive a potential Lyme disease vaccine. Multivariable multinomial logistic regression models were used to quantify the association of sociodemographic characteristics and Lyme disease vaccine attitudes with willingness to be vaccinated.

Results: Surveys were completed by 3313 respondents (6% response rate). We estimated that 64% of residents were willing to receive a Lyme disease vaccine, while 30% were uncertain and 7% were unwilling. Compared to those who were willing, those who were uncertain were more likely to be parents, adults 45–65 years old, non-White, have less than a bachelor's degree, or have safety concerns about a potential Lyme disease vaccine. Those who were unwilling were also more likely to be non-White, have less than a bachelor's degree, or have safety concerns about a potential Lyme disease vaccine. In addition, the unwilling had low confidence in vaccines in general, had low perceived risk of contracting Lyme disease, and said they would not be influenced by a positive recommendation from a healthcare provider.

Discussion: Overall, willingness to receive a Lyme disease vaccine was high. Effective communication by clinicians regarding safety and other vaccine parameters to those groups who are uncertain will be critical for increasing vaccine uptake and reducing Lyme disease incidence.

© 2021 Published by Elsevier Ltd.

1. Background

Lyme disease (LD) is a multi-system illness caused by infection with *Borrelia burgdorferi*. These spirochetes are transmitted to humans and animals by the bite of infected *Ixodes* species ticks, primarily in northeastern, mid-Atlantic, and upper-midwestern regions of the United States (US) [1,2]. Incidence has been increasing, with over 30,000 cases reported annually to the Centers for Disease Control and Prevention (CDC) [1]. However, recent studies

* Corresponding author.

E-mail addresses: vhx8@cdc.gov (S.A. Hook), amberjean.hansen@yale.edu (AmberJean P. Hansen), sara.niesobecki@yale.edu (S.A. Niesobecki), james.meek@yale.edu (J.I. Meek), jenna.bjork@state.mn.us (J.K.H. Bjork), erin.kough@state.mn.us (E.M. Kough), molly.peterson@state.mn.us (M.S. Peterson), elizabeth.schiffman@state.mn.us (E.K. Schiffman), heather.rutz@maryland.gov (H.J. Rutz), adam.rowe@health.ny.gov (A.J. Rowe), jennifer.white@health.ny.gov (J.L. White), jennifer.peel@colostate.edu (J.L. Peel), bkb5@cdc.gov (B.J. Biggerstaff), cue0@cdc.gov (A.F. Hinckley).

<https://doi.org/10.1016/j.vaccine.2021.11.065>
0264-410X/© 2021 Published by Elsevier Ltd.

have estimated that there are nearly 500,000 diagnosed cases annually [3–5]. Early symptoms of LD most often include a characteristic bull's-eye rash known as erythema migrans, as well as flu-like symptoms [6]. If left untreated, the disease can disseminate to cause more severe manifestations, such as arthritis, meningitis, or carditis, the last of which can be fatal in rare cases. Most patients will experience a full recovery after antibiotic treatment, although some may continue to experience symptoms related to disease sequelae [6–10]. Despite the availability of antibiotic treatment, an effective LD vaccine is needed to prevent severe outcomes and long-term symptoms and thereby reduce fiscal burdens on patients and healthcare systems. Further, currently available personal and yard-based prevention methods have not been sufficient to stem rising case numbers, highlighting the need for a population-level prevention modality such as a vaccine [11,12 13].

A safe and efficacious vaccine for LD called LYMERix was available for persons aged 15–70 years from 1998 until 2002 in the US [14,15]. This vaccine conferred protection based on a recombinant outer surface protein A (rOspA) of *B. burgdorferi*. In 2002, it was voluntarily discontinued by the manufacturer, reportedly due to poor sales [16]. However, several factors have been highlighted as reasons contributing to low demand. Most importantly, it was not available for children under 15 years, one of the highest risk age groups. Further, some have cited tepid and cumbersome recommendations by the Advisory Committee on Immunization Practices (ACIP) as a potential reason for low demand by clinicians and the public [17,18]. Vocal opposition by some Lyme disease patient advocacy groups, based on unsubstantiated claims that the vaccine caused Lyme arthritis, is also thought to have played a role in LYMERix's withdrawal [19–22]. The introduction and withdrawal of LYMERix also inauspiciously coincided with the then nascent anti-vaccination movement [22]. Since its withdrawal, the number of LD cases reported annually has nearly doubled.

After nearly two decades without an LD vaccine, new candidates are in development, with initial results showing favorable safety and immunogenicity profiles and potential availability in the next several years [23–25]. While rising LD incidence would ostensibly result in increased demand for a vaccine, the controversial climate surrounding LD [26] and general vaccine hesitancy among some groups [27–30] necessitate further investigation of the acceptability of a LD vaccine among potential consumers. The primary objective of this study was to estimate what proportion of people living in states with a high incidence of LD would be willing to receive a new LD vaccine. The secondary objective was to evaluate factors associated with willingness to receive a LD vaccine.

2. Methods

2.1. Study design and sampling

In the summer of 2018, we conducted a population-based, cross-sectional survey using address-based sampling of persons living in four states with high incidence of LD [31]. The target population included all residents of Connecticut, Maryland, Minnesota, and New York, excluding New York City due to low incidence of LD. The sampling frame included all households with residential addresses listed in the U.S. Postal Service (USPS) database in these areas. We used a stratified, two-stage sampling design where the strata were counties from the above-mentioned states. The primary sampling unit was the household, while the unit of observation was the individual, with a single individual selected within the household. Addresses were purchased from a marketing company that receives updated information on a monthly basis directly from USPS based on change of address submissions. Household

addresses were stratified according to county, and the number of addresses selected per county was allocated proportional to county population size. Households were randomly selected within counties. An individual within the household was selected as the one who had the most recent birthday, regardless of age, an established technique to approximate random sampling [32]. For minors selected, parents or guardians ≥ 18 years of age provided responses; responses from those < 18 years were excluded. Subsequently, the term “respondent” will refer to those about whom information was collected.

To estimate the proportion of residents willing to receive a LD vaccine, the sample size calculation parameters included a conservative estimate of 50% of participants responding “Yes” for willingness to receive a LD vaccine; $\alpha = 0.01$; an acceptable margin of error of $\pm 5\%$; and 2 clusters for multi-stage sampling [33]. These parameters resulted in a required sample size of 665 respondents per state (2660 respondents total). Based on a 2016 survey using address-based sampling in Connecticut and Maryland [34], we anticipated a 5% response rate and, therefore, recruited 13,300 individuals per state (53,200 total) to obtain a sample representative of the populations in these states (including responses for both adults and children), in the absence of non-response.

2.2. Data collection

Recruitment, enrollment, and survey completion occurred during June–July 2018, with data collection corresponding with peak tickborne disease activity. An invitation postcard explaining the survey in English was mailed to each randomly selected household. The postcard provided a web link, quick response (QR) barcode, and a unique access code to complete the online survey; alternatively, respondents could choose to complete the survey over the phone with study coordinators. A reminder was mailed two weeks later, and the online surveys were open for approximately four weeks.

Sociodemographic information was collected from respondents (Table 1). An additional variable for metropolitan status (large central metropolitan area vs other) by county was created using the urban–rural classification scheme from the National Center for Health Statistics [35]. The main outcome variable was whether the respondent would be willing to receive a LD vaccine if one were available (or vaccinate the minor, if a parent respondent). The following covariates were also collected from survey responses (Appendix, Table A1): LD vaccine safety concerns; vaccine cost concerns; acceptance of vaccine recommendations from a healthcare provider (HCP); history of LD diagnosis among household members; level of concern about getting LD; time spent in tick habitat; whether vaccines, in general, benefit people; primary source for LD information; and primary location for receiving vaccinations.

2.3. Analysis

The data were weighted to account for the unequal selection probabilities per respondent for the two-stage sampling design [32,36]. We compared the sample distributions of age and gender to known population totals using chi-squared goodness-of-fit tests, and as necessary, conducted post-stratification according to county population distributions of age and gender to reduce sampling error and nonresponse error [36–39]. All analyses were conducted using the weighted, post-stratified dataset, and all analyses incorporated the sampling design into standard error and confidence interval computation and statements of inference. We also evaluated non-random missingness in our outcome variable related to non-response (*i.e.*, selection bias) using Heckman-type selection models [40–42] (Appendix, Section 1 and Table A4).

Table 1
Respondent characteristics and willingness to receive a Lyme disease (LD) vaccine, weighted % (95% CI).

Characteristic	All**	Willingness to receive a LD vaccine*		
		Yes	No	Don't Know
Total, N = 3206		64 (62, 65), n = 2098	7 (6, 8), n = 190	30 (29, 31), n = 918
<i>Demographics</i>				
Gender ***				
Female	54	54 (53, 54)	64 (59, 70)	54 (52, 56)
Male	46	46 (46, 47)	36 (30, 41)	46 (44, 48)
Age category *** (years)				
<18	15	14 (13, 15)	13 (10, 16)	19 (17, 20)
18–44	33	36 (35, 37)	34 (29, 40)	28 (26, 30)
45–64	34	32 (31, 33)	39 (33, 45)	38 (36, 40)
65+	17	18 (18, 19)	14 (11, 17)	16 (14, 17)
State				
CT	20	21 (21, 22)	17 (12, 21)	16 (15, 18)
MD	27	27 (26, 28)	24 (19, 29)	29 (27, 31)
MN	20	20 (19, 21)	24 (20, 28)	18 (17, 20)
NY	33	32 (31, 33)	35 (29, 41)	37 (35, 39)
Race				
White	85 (84, 86)	87 (86, 88)	75 (69, 81)	81 (79, 83)
Non-White	15 (14, 16)	13 (12, 14)	25 (19, 31)	19 (17, 21)
Education				
Some college or less	35 (33, 36)	31 (29, 32)	50 (44, 56)	39 (37, 41)
Bachelor's degree or higher	65 (64, 67)	69 (68, 71)	50 (44, 56)	61 (59, 63)
Metropolitan status				
Large central metro area	28	28 (27, 29)	38 (33, 44)	26 (24, 28)
Other	72	72 (71, 73)	62 (56, 67)	74 (72, 76)
<i>LD history, attitudes, and practices</i>				
Past LD diagnosis in household				
Yes	18 (17, 19)	21 (20, 22)	14 (9, 19)	13 (12, 15)
No	82 (81, 83)	79 (78, 80)	86 (81, 91)	87 (85, 88)
Concern about future LD diagnosis				
Yes	86 (85, 86)	94 (93, 95)	56 (50, 62)	74 (72, 76)
No	14 (14, 15)	6 (5, 7)	44 (38, 50)	26 (24, 28)
Time spent in tick habitat				
At least weekly	71 (70, 73)	82 (80, 83)	51 (45, 57)	54 (52, 56)
Monthly or less	29 (27, 30)	18 (17, 20)	49 (43, 55)	46 (44, 48)
Current use of LD prevention measures				
Yes	92 (91, 93)	94 (93, 95)	87 (83, 91)	90 (88, 91)
No	8 (7, 9)	6 (5, 7)	13 (9, 17)	10 (9, 12)
Confidence in LD prevention measures				
Yes	70 (68, 71)	67 (65, 68)	81 (76, 85)	74 (71, 76)
No	30 (29, 32)	33 (32, 35)	19 (15, 24)	26 (24, 29)
Confidence in general vaccines				
Yes	94 (93, 95)	98 (97, 98)	69 (64, 74)	91 (89, 93)
No	6 (5, 7)	2 (2, 3)	31 (26, 36)	9 (7, 11)
<i>LD vaccine attitudes</i>				
LD vaccine safety concerns				
Yes	71 (70, 72)	68 (66, 69)	80 (75, 84)	75 (74, 77)
No	29 (28, 30)	32 (31, 34)	20 (16, 25)	25 (23, 26)
Healthcare provider influence on LD vaccination				
Yes	89 (88, 89)	93 (92, 94)	57 (52, 63)	87 (85, 89)
No	11 (11, 12)	7 (6, 8)	43 (37, 48)	13 (11, 15)
LD vaccine cost concerns				
Yes	63 (62, 65)	66 (65, 68)	34 (28, 40)	64 (62, 67)
No	37 (35, 38)	34 (32, 35)	66 (60, 72)	36 (33, 38)

* All comparisons made in stratified analyses using Pearson chi-squared tests with Rao and Scott design-based adjustments had resultant p values ≤ 0.001 .

** County distributions of gender and age were used for post-stratification; as such, these point estimates for the overall sample are fixed at the population values and have no associated interval estimate. Because state and metropolitan status are based on county population totals, these point estimates are also fixed.

*** Gender and age categories represent the potential vaccinee, i.e., adult respondents and the children for whom parents responded.

To estimate the proportion of people in Connecticut, Maryland, Minnesota, and New York willing to receive a LD vaccine, summary statistics were computed for the three-level response for willingness to receive a vaccine. Additionally, descriptive analyses were conducted for the following independent variables: sociodemographic characteristics; LD history, attitudes, and practices; vaccine attitudes; primary sources of LD information; and primary location for receiving vaccines.

To evaluate factors associated with willingness to receive a LD vaccine, we cross-classified the outcome with the above mentioned independent variables, and Pearson chi-squared tests with Rao and Scott design-based adjustments were used to evaluate differences in the outcome across levels of each independent variable [43]. Because our outcome of willingness to vaccinate had three, unordered levels, multivariable multinomial logistic regression models were used to quantify the association between LD vaccina-

Table 2
Unadjusted and adjusted odds ratios for Lyme disease (LD) vaccination responses using multinomial logistic regression.

Variable	LD vaccination responses (ref. = Yes)			
	No OR (95% CI)	aOR (95% CI)	Don't Know OR (95% CI)	aOR (95% CI)
Vaccinee age category^a (ref. = 65 + years)				
<18	1.19 (0.38, 3.68)	NA	1.60 (1.06, 2.42)	NA
18–44	1.23 (0.69, 2.19)	NA	0.91 (0.66, 1.26)	NA
45–64	1.59 (0.88, 2.85)	NA	1.40 (1.07, 1.85)	NA
Gender^a (ref. = Male)				
Female	1.55 (0.90, 2.68)	NA	1.00 (0.77, 1.31)	NA
State¹ (ref. = Connecticut)				
Maryland	1.13 (0.60, 2.13)	1.16 (0.61, 2.19)	1.40 (0.99, 1.98)	1.42 (1.01, 1.99)
Minnesota	1.54 (0.77, 3.07)	1.51 (0.76, 3.00)	1.20 (0.83, 1.73)	1.19 (0.82, 1.73)
New York	1.41 (0.79, 2.50)	1.41 (0.80, 2.48)	1.52 (1.05, 2.20)	1.52 (1.05, 2.19)
Race² (ref. = White)				
Non-White	2.24 (1.18, 4.26)	2.29 (1.21, 4.32)	1.55 (1.10, 2.18)	1.54 (1.10, 2.17)
Education³ (ref. = ≥ Bachelor's degree)				
<Bachelor's degree	2.29 (1.35, 3.88)	2.21 (1.28, 3.83)	1.45 (1.12, 1.87)	1.47 (1.13, 1.91)
LD vaccine safety concerns⁴ (ref. = No)				
Yes	1.86 (1.12, 3.1)	2.62 (1.49, 4.60)	1.48 (1.07, 2.03)	1.99 (1.42, 2.78)
HCP influence on LD vaccination⁵ (ref. = Yes)				
No	9.32 (5.43, 16.01)	5.21 (2.72, 10.00)	1.92 (1.30, 2.84)	1.42 (0.94, 2.15)
LD vaccine cost concerns⁶ (ref. = No)				
Yes	0.26 (0.16, 0.43)	0.36 (0.20, 0.64)	0.92 (0.74, 1.16)	1.07 (0.82, 1.39)

^a Unadjusted models only; no potential confounders were included in these models.

¹ State model adjusted for age category and education.

² Race model adjusted for metro status.

³ Education model adjusted for age category, gender, state, race, and metro status.

⁴ LD vaccine safety concerns model adjusted for age category, gender, education, HCP recommendation, past LD diagnosis in household, concern about future LD diagnosis, time spent in tick habitat, current use of LD prevention measures, general confidence in vaccines.

⁵ HCP (healthcare provider) influence on LD vaccination model adjusted for age category, gender, education, past LD diagnosis in household, concern about future LD diagnosis, time spent in tick habitat, general confidence in vaccines.

⁶ LD vaccine cost concerns model adjusted for age category, gender, state, education, HCP recommendation, concern about future LD diagnosis, time spent in tick habitat, current use of LD prevention measures, general confidence in vaccines.

tion responses and independent variables of interest. For each model, we used “Yes” responses to willingness to be vaccinated as the reference group to which “No” and “Don't know” responses were compared. The independent variables of interest included sociodemographic characteristics (*i.e.*, gender, age category, state, race, and education), LD vaccine safety concerns, LD vaccine cost concerns, and positive recommendation for the LD vaccine from an HCP. The last three independent variables were dichotomized for analysis (Yes = “Some” or “A lot”; No = “Not at all” or “Don't know”). Separate models were built for each independent variable of interest with a specific set of potential confounders identified *a priori* (Appendix, Table A2), and model diagnostics were conducted for each model fit. Multinomial logistic regression model results are presented as unadjusted and adjusted odds ratios (ORs) and 95% confidence intervals (CIs).

Survey development, administration, data collection, and data management were conducted using the Research Electronic Data Capture (REDCap) software hosted at Yale University [44,45]. R version 3.5.2 [46–51] was used for all analyses. This study was conducted through TickNET, a public health network composed of researchers at state health departments, universities, and CDC who collaborate on tickborne disease research and surveillance [52]. Research approval and waiver of documentation of informed consent were obtained from institutional review boards at CDC, Connecticut Department of Public Health, Maryland Department of Health, Minnesota Department of Health, New York State Department of Health, and Yale University. Respondents' participation in the survey indicated consent.

3. Results

The survey response rate was 6% ($n = 3313$). Fifty-nine records were ineligible due to the following: missing age data ($n = 38$), the

respondent not being the person in the household with the last birthday ($n = 15$), the adult respondent not being the one to make vaccination decisions for the selected minor ($n = 1$), or the respondent not answering the main outcome question regarding willingness to be vaccinated ($n = 5$). An additional 48 records with missing gender information were removed prior to analysis because gender information was necessary for post-stratification. The resulting sample available for analysis was 3206 records. The coefficient of the inverse Mill's ratio resulting from Heckman selection models indicated no evidence of significant selection bias (Appendix, Table A4).

Individuals in the sample were older with a higher proportion female compared to the source population; therefore, we post-stratified the data on age and gender as described above [36–39]. The following proportions of demographic characteristics were fixed by post-stratification: 54% of residents were female, 17% were aged ≥ 65 years, 33% were from New York, and 28% lived in a large central metropolitan area (Table 1). In weighted analysis, we estimated that 15% of residents were parents, 85% were White, and 65% had a bachelor's degree or higher; CIs are reported in Table 1.

For our outcome of interest, we estimated that 64% ($n = 2098$) of residents were willing to receive a LD vaccine, while 7% ($n = 190$) were not willing and 30% ($n = 918$) were uncertain (Table 1). We estimated that 18% of residents experienced a past LD diagnosis in their household, and 86% expressed concern about a future LD diagnosis. An estimated 71% of residents spent time in tick habitat at least weekly. Nearly all residents (92%) used some type of LD prevention measure, while 70% were confident that available measures can prevent LD. The vast majority (94%) were confident that recommended vaccines benefit people. Regarding LD vaccine attitudes, the majority of residents had concerns about vaccine safety (71%) and cost (63%), and the majority (89%) indicated that a pos-

itive recommendation from an HCP for the LD vaccine would influence their willingness to be vaccinated. In stratified analyses, differences in willingness to be vaccinated were observed for all characteristics and were significant at $\alpha = 0.05$.

Overall, we estimated that the top sources of LD information for residents were health websites (29%, 95% CI: 28%, 30%), search engines (22%, 95% CI: 21%, 23%), and HCPs (21%, 95% CI: 20%, 22%) (Fig. 1), with similar proportions for those who said “Yes” and “Don’t know” to potential LD vaccination (Appendix, Fig. A1). Among those who said “No” to potential LD vaccination, a lower proportion (22%, 95% CI: 17%, 27%) cited health websites as a primary source of LD information, and a higher proportion cited search engines (25%, 95% CI: 19%, 31%) and social media (6%, 95% CI: 2%, 9%), compared to residents overall and those who said “Yes” and “Don’t know” to LD vaccination (Appendix, Fig. A1).

Overall, the top three locations for receiving vaccinations were HCP offices, clinics, or hospitals (82%, 95% CI: 81%, 83%); pharmacies (12%, 95% CI: 11%, 12%); and workplaces (3%, 95% CI: 2%, 3%) (Fig. 2). Proportions were similar for those who said “Yes” and “Don’t know” to potential LD vaccination, while a higher proportion of those who said “No” said they “do not get vaccines” (14%, 95% CI: 10%, 18%) or that they “Don’t know” their primary location for receiving vaccination (5%, 95% CI: 4%, 6%), compared to residents overall and those who said “Yes” and “Don’t know” to LD vaccination (Appendix, Fig. A2).

Table 2 shows the estimated unadjusted and adjusted ORs and 95% CIs resulting from survey-weighted, multivariable, multinomial logistic regression analysis. In terms of sociodemographic characteristics, the odds of parents of minors responding “Don’t know” (vs. “Yes”) to LD vaccination was 1.6 times that of the reference group, those 65 years and older (OR: 1.60, 95% CI: 1.06, 2.42). The odds of those aged 45–64 years responding “Don’t know” were also higher compared to those 65 years and older (OR: 1.40, 95% CI: 1.07, 1.85). Females had only slightly higher odds of responding “No” (vs. “Yes”) to LD vaccination compared to males (OR: 1.55, 95% CI: 0.90, 2.68) and did not have higher odds of responding “Don’t know”. Those in Maryland and New York had higher odds of responding “Don’t know” to LD vaccination compared to those

in Connecticut (aOR: 1.42, 95% CI: 1.01, 1.99 and aOR: 1.52, 95% CI: 1.05, 2.19, respectively). No differences were found among states for “No” responses. Non-White residents had higher odds of responding “No” to LD vaccination (aOR: 2.29, 95% CI: 1.21, 4.32) and “Don’t know” (aOR: 1.54, 95% CI: 1.10, 2.17) compared to White residents. Those with less than a bachelor’s degree had higher odds of responding “No” (aOR: 2.21, 95% CI: 1.28, 3.83) and “Don’t know” (aOR: 1.47, 95% CI: 1.13, 1.91) to LD vaccination compared to those with more education.

In terms of attitudes toward a LD vaccine, those with safety concerns had higher odds of responding “No” and “Don’t know” to LD vaccination (aOR: 2.62, 95% CI: 1.49, 4.6; aOR: 1.99, 95% CI: 1.42, 2.78, respectively) compared to those without safety concerns. Those who said HCP recommendation would not influence their willingness to be vaccinated had much higher odds of responding “No” (aOR: 5.21, 95% CI: 2.72, 10.00) but only slightly higher odds of responding “Don’t know” (aOR: 1.42, 95% CI: 0.94, 2.15). Finally, those with LD vaccine cost concerns had lower odds of responding “No” to LD vaccination (aOR: 0.36, 95% CI: 0.20, 0.64) compared to those without cost concerns.

4. Discussion

We estimate that over 60% of residents in states with a high incidence of Lyme disease would be willing to receive a LD vaccine if one were available. Approximately 30% of residents were unsure about vaccination, and they were more likely to be parents making decisions for their children, adults 45–65 years of age, non-White, have less than a bachelor’s degree, or have concerns about the safety of a potential LD vaccine. Targeted vaccine communications by HCPs to these groups, especially those in the age groups at highest risk for LD (children aged 5–10 years and adults aged 45–55 years [1]) may increase uptake of a LD vaccine. Less than 10% of residents indicated that they were not willing to be vaccinated. They were also more likely to be non-White, have less than a bachelor’s degree, or have concerns about the safety of a potential LD vaccine, but they also would not be influenced by a positive recommendation from a HCP, have low confidence in vaccines in general,




Fig. 1. Residents' primary source for LD information. The gray bars represent the weighted percent. The black lines represent 95% confidence intervals.




Fig. 2. Residents' primary location for receiving vaccination. The gray bars represent the weighted percent. The black lines represent 95% confidence intervals. "HCP" denotes healthcare provider.

and have low perceived risk of contracting LD. Targeted outreach may be unlikely to change these groups' willingness to receive a LD vaccine. Alternatively, these groups may have low perceived risk of LD because of truly being at low risk (e.g., those who do not spend time outdoors in tick habitat), and they may not benefit from LD vaccination.

A 2002 study among parents in Nassau County, New York evaluated whether parents would request the LYMERix vaccine for their children, if and when it became available. The vast majority said they would "definitely" (23%) or "likely" (65%) request it, followed by those "unlikely" (9%) to request it and those who would not (3%) [53]. While this response scale differs from that in our study, these results are similar to ours, with the majority willing to be vaccinated and few declining. Another study evaluated a LYMERix vaccination program among New York State Department of Health employees at risk for occupational tick exposure. While only 16% of employees chose to be vaccinated, the majority of non-recipients reported safety as a major concern, as seen in our results [54].

Prior to this study there has been little research on acceptability of a potential new LD vaccine, though a 2016 convenience sample survey conducted in Connecticut and Maryland counties with a high incidence of LD found that the majority of respondents were likely to receive a potential LD vaccine, with 49% "very likely", 35% "somewhat likely", 8% "somewhat unlikely", and 7% "very unlikely" [34]. Similarly, a nationwide population-based survey conducted in 2014 and 2015 found that 65% of respondents in high incidence states would be "likely" to receive a potential LD vaccine [55]. Additionally, a qualitative research study conducted in 2018 using focus groups of those at high risk for LD showed that 57% would be "very likely" to receive a LD vaccine [69]. Again, while the response scales of these studies differ from the present study, our estimates of potential vaccine uptake are concordant.

Demographic disparities in vaccination coverage are common and complex for both compulsory childhood vaccines and for recommended, non-compulsory vaccines for adults and children

[27,28,56–59]. Our finding that those who are non-White or those with lower education are more likely to respond "No" and "Don't know" to LD vaccination contrasts somewhat with studies on childhood vaccines. In Arizona, non-medical exemption rates (i.e., vaccine refusals) among kindergarteners were higher in schools with a higher proportion of White children and a lower proportion of free lunches (as a proxy for income) [27]. In a nationwide survey, more White parents reported being unsure about or refusing childhood vaccinations versus other racial groups [28]. However, another nationwide survey found demographic differences when comparing under-vaccinated children with unvaccinated children; under-vaccinated children tended to be Black, have a mother without a college degree, and have lower household income while unvaccinated children tended to be White, have a mother with a college degree, and have higher household income [56]. However, our survey was not parent-specific, and our sample includes only a small proportion of parent respondents. Our results for a voluntary LD vaccine are likely more comparable to annual reports of coverage for recommended, non-compulsory vaccines for adults. Annually, these reports show higher coverage generally for White persons compared with most other racial groups [57,58]. People from racial and ethnic minority groups are at risk for LD, and vaccine communications should focus on these groups in endemic areas.

Vaccine safety concerns are often cited as reasons for delaying or refusing vaccinations generally among both parents and adults, and these concerns were also an important factor in LYMERix vaccination decisions, despite it being proven to be safe [28–30,53,54,56,60,61]. Our results show that safety will also be an important consideration in future LD vaccination decisions. A new LD vaccine may spawn additional safety concerns given that the waning demand for LYMERix was due, in part, to safety concerns, albeit unfounded. However, current vaccine candidates do not include the alleged, arthritis-causing epitope present in the LYMERix vaccine, which may assuage concerns for some [62]. Further, many studies, including this one, have shown that a positive

recommendation for vaccination from an HCP has a significant influence on the vaccination decision and can increase uptake [61,63,64]. While other factors such as efficacy, convenience, and LD risk, among others, will undoubtedly play a role in uptake of a potential LD vaccine, effectively communicating its safety profile will be critical, and HCPs should be primary communicators of this information to the public [65]. As such, it will be important for public health practitioners to work with HCPs to develop messaging and other tools for discussing LD vaccines with patients.

These results must be interpreted in the context of several potential limitations. While we anticipated and accounted for a low response rate in our sample size calculations, such large non-response may affect the validity of our estimate of vaccine uptake due to non-response error. For example, it is possible that those who do not perceive themselves to be at risk for LD had low interest in the survey and chose not to respond. These non-respondents may be likely to decline LD vaccination due to their perceived low risk, thereby causing an overestimate of the proportion willing to be vaccinated in our sample, compared to the target population. Similarly, those who perceive themselves to be at high risk for LD may have had keen interest in participating in the survey, and their perceived risk may lead to higher willingness to be vaccinated, again leading to an overestimate of vaccine acceptance. However, post-stratification was intended to mitigate this non-response error, and Heckman selection model results did not reveal significant selection bias. Further, our results are very similar to other LD vaccine acceptability studies with different sampling methods, as noted above. In terms of information bias, most survey questions, including the willingness to vaccinate outcome and independent variables of interest, concerned respondents' opinions, making recall or misclassification error unlikely. However, given the hypothetical nature of the survey questions, the estimate of intention to receive a LD vaccine may change as more information on vaccine parameters becomes available or may differ from actual vaccine uptake. For example, results were mixed for studies evaluating the correlation between intention to receive a vaccine and actual uptake during the 2009–2010 influenza A/H1N1 pandemic in the United States [66–68]. Lastly, while our results are generalizable to the populations of participating states, excluding residents of New York City, these results may or may not be generalizable to other states with a high incidence of LD. However, the states in this study represent a range of endemicity, from fully endemic in Connecticut to focally endemic in parts of Minnesota and New York; therefore, results are likely applicable to other endemic states, such as Massachusetts and Wisconsin, but may not apply to states where LD is emerging, such as Michigan and West Virginia.

In anticipation of a new LD vaccine coming to market, future studies should further evaluate parent-specific vaccine concerns, given that children are at high risk for LD and may benefit most from the vaccine. Additional evaluations of vaccine acceptability will also be needed once safety, efficacy, dosing, and immunogenicity data is available for a new vaccine. Our estimate of potential vaccine uptake provides important information for ACIP recommendations and may be used in economic evaluations of a potential vaccine. Lastly, our characterization of the factors affecting willingness to receive a potential LD vaccine can inform future communication and education efforts with clinicians and the public to increase awareness and uptake of a vaccine.

5. Conclusions

LD incidence is increasing, despite current prevention options. A new LD vaccine could substantially reduce disease incidence if vaccine uptake is high. The majority of residents in four high incidence

states would be willing to receive a LD vaccine if one were available. Effective communication by clinicians regarding safety, efficacy, and other vaccine parameters to those demographic groups who are uncertain about LD vaccination will be critical for increasing vaccine uptake and reducing LD incidence.

Funding

This work was supported by the Centers for Disease Control and Prevention.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors would like to thank Stacy Greby, Allison Fisher, Carla Black, Amy Schwartz, Marc Fischer, and David Neitzel for their valuable contributions to the design of this survey. We would also like to thank Paul Mead, Sheryl Magzamen, Jude Bayham, and Brooke Anderson for their review of the manuscript.

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.vaccine.2021.11.065>.

References

- [1] Schwartz AM, Hinckley AF, Mead PS, Hook SA, Kugeler KJ. Surveillance for Lyme disease – United States, 2008–2015. *MMWR Surveill Summ* 2017;66(22):1–12.
- [2] Rosenberg R, Lindsey NP, Fischer M, Gregory CJ, Hinckley AF, Mead PS, et al. Vital signs: trends in reported vectorborne disease cases – United States and Territories, 2004–2016. *MMWR Morb Mortal Wkly Rep* 2018;67(17):496–501.
- [3] Nelson CA, Saha S, Kugeler KJ, Delorey MJ, Shankar MB, Hinckley AF, et al. Incidence of clinician-diagnosed Lyme disease, United States, 2005–2010. *Emerg Infect Dis* 2015;21(9):1625–31.
- [4] Hinckley AF, Connally NP, Meek JI, Johnson BJ, Kemperman MM, Feldman KA, et al. Lyme disease testing by large commercial laboratories in the United States. *Clin Infect Dis* 2014;59:676–81.
- [5] Kugeler KJ, Schwartz AM, Delorey MJ, Mead PS, Hinckley AF. Estimating the frequency of Lyme disease diagnoses, United States, 2010–2018. *Emerg Infect Dis* 2021;27(2):616–9.
- [6] Wormser GP, Dattwyler RJ, Shapiro ED, Halperin JJ, Steere AC, Klempner MS, et al. The clinical assessment, treatment, and prevention of Lyme disease, human granulocytic anaplasmosis, and babesiosis: clinical practice guidelines by the Infectious Diseases Society of America. *Clin Infect Dis* 2006;43:1089–134.
- [7] Magid D, Schwartz B, Craft J, Schwartz JS. Prevention of Lyme disease after tick bites. A cost-effectiveness analysis. *N Engl J Med* 1992;327(8):534–41.
- [8] Nichol G, Dennis DT, Steere AC, Lightfoot R, Wells G, Shea B, et al. Test-treatment strategies for patients suspected of having Lyme disease: a cost-effectiveness analysis. *Ann Intern Med* 1998;128:37–48.
- [9] Meltzer MI, Dennis DT, Orloski KA. The cost effectiveness of vaccinating against Lyme disease. *Emerg Infect Dis* 1999;5(3):321–8.
- [10] Weitzner E, McKenna D, Nowakowski J, Scavarda C, Dornbush R, Bittker S, et al. Long-term assessment of post-treatment symptoms in patients with culture-confirmed early Lyme disease. *Clin Infect Dis* 2015;61(12):1800–6.
- [11] CDC. Lyme disease charts and figures: historical data; 2019.
- [12] Hayes EB, Piesman J. How can we prevent Lyme disease? *N Engl J Med* 2003;348(24):2424–30.
- [13] Gomes-Solecki M, Arnaboldi PM, Backenson PB, Benach JL, Cooper CL, Dattwyler RJ, et al. Protective immunity and new vaccines for Lyme disease. *Clin Infect Dis* 2020;70(8):1768–73.
- [14] Poland GA, Jacobson RM. The prevention of Lyme disease with vaccine. *Vaccine* 2001;19(17–19):2303–8.
- [15] ACIP. Recommendations for the use of Lyme disease vaccine recommendations of the Advisory Committee on Immunization Practices (ACIP). *Morb Mortal Wkly Rep* 1999;48:1–17.
- [16] Shen AK, Mead PS, Beard CB. The Lyme disease vaccine—a public health perspective. *Clin Infect Dis* 2011;52(Suppl 3):S247–52.

- [17] Plotkin SA. Correcting a public health fiasco: the need for a new vaccine against Lyme disease. *Clin Infect Dis* 2011;52(Suppl 3):s271–5.
- [18] Plotkin SA. Need for a new Lyme disease vaccine. *N Engl J Med* 2016;375(10):911–3.
- [19] Aronowitz RA. The rise and fall of the Lyme disease vaccines: a cautionary tale for risk interventions in American medicine and public health. *Milbank Q* 2012;90:250–77.
- [20] Lathrop SL, Ball R, Haber P, Mootrey GT, Braun MM, Shadomy SV, et al. Adverse event reports following vaccination for Lyme disease: December 1998–July 2000. *Vaccine* 2002;20(11–12):1603–8.
- [21] Ball R, Shadomy SV, Meyer A, Huber BT, Leffell MS, Zachary A, et al. HLA type and immune response to *Borrelia burgdorferi* outer surface protein a in people in whom arthritis developed after Lyme disease vaccination. *Arthritis Rheum* 2009;60(4):1179–86.
- [22] Nigrovic LE, Thompson KM. The Lyme vaccine: a cautionary tale. *Epidemiol Infect* 2007;135(1):1–8.
- [23] Flaherty B. Can a new Lyme disease vaccine overcome a history of distrust and failure? *STAT*; 2019.
- [24] Wang Y, Esquivel R, Flingai S, Schiller ZA, Kern A, Agarwal S, et al. Anti-OspA DNA-encoded monoclonal antibody prevents transmission of spirochetes in tick challenge providing sterilizing immunity in mice. *J Infect Dis* 2019;219:1146–50.
- [25] Wang Y, Kern A, Boatright NK, Schiller ZA, Sadowski A, Ejemel M, et al. Pre-exposure prophylaxis with OspA-specific human monoclonal antibodies protects mice against tick transmission of Lyme disease spirochetes. *J Infect Dis* 2016;214(2):205–11.
- [26] Lantos PM. Chronic Lyme disease: the controversies and the science. *Expert Rev Anti Infect Ther* 2011;9(7):787–97.
- [27] Birnbaum MS, Jacobs ET, Ralston-King J, Ernst KC. Correlates of high vaccination exemption rates among kindergartens. *Vaccine* 2013;31(5):750–6.
- [28] Gust DA, Darling N, Kennedy A, Schwartz B. Parents with doubts about vaccines: which vaccines and reasons why. *Pediatrics* 2008;122(4):718–25.
- [29] Lutz CS, Fink RV, Cloud AJ, Stevenson J, Kim D, Fiebelkorn AP. Factors associated with perceptions of influenza vaccine safety and effectiveness among adults, United States, 2017–2018. *Vaccine* 2020;38(6):1393–401.
- [30] Salmon DA, Dudley MZ, Glanz JM, Omer SB. Vaccine hesitancy: causes, consequences, and a call to action. *Am J Prev Med* 2015;49(6):S391–8.
- [31] Link MW, Battaglia MP, Frankel MR, Osborn L, Mokdad AH. A Comparison of address-based sampling (ABS) versus random-digit dialing (RDD) for general population surveys. *Publ Opin Quart* 2008;72(1):6–27.
- [32] Fowler FJ. *Survey research methods*. 5th ed. Los Angeles: SAGE; 2014.
- [33] CDC. *Epi Info*. In: CDC, editor. 7.2.2.2 ed 2017.
- [34] Niesobecki S, Hansen AJ, Rutz H, Mehta S, Feldman K, Meek J, et al. Knowledge, attitudes, and behaviors regarding tick-borne disease prevention in endemic areas. *Ticks Tick Borne Dis* 2019;10(6).
- [35] Ingram DD, Franco SJ. NCHS urban-rural classification scheme for counties. *Vital Health Stat* 2013;2(2014):1–73.
- [36] Lumley T. Analysis of complex survey samples. *J Stat Softw* 2004;9:1–19.
- [37] Lumley T. *Complex surveys: a guide to analysis using R*. Hoboken, New Jersey: John Wiley & Sons, Inc.; 2010.
- [38] Sampling LSL. *Design and analysis (advanced series)*. 2nd ed. Boston, MA: Brooks/Cole; 2010.
- [39] Sarndal C-E, Lundstrom S. Estimation in surveys with nonresponse. West Sussex PO19 8SQ, England: John Wiley & Sons, Ltd; 2005.
- [40] Heckman JJ. Sample selection bias as a specification error. *Econometrica* 1979;47(1):153. <https://doi.org/10.2307/1912352>.
- [41] Amemiya T. Tobit models: a survey. *J Econ* 1984;24(1–2):3–61.
- [42] Toomet O, Henningsen A. Sample selection models in R: package *sampleselection*.
- [43] Rao J, Scott J. On Chi-squared tests for multiway contingency tables with cell proportions estimated from survey data. *Ann Stat* 1984;12:46–60.
- [44] Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L, et al. The REDCap consortium: building an international community of software platform partners. *J Biomed Inform* 2019;95:103208. <https://doi.org/10.1016/j.jbi.2019.103208>.
- [45] Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;42(2):377–81.
- [46] Wickham H. tidyverse: easily install and load the 'Tidyverse'. R package version 1.2.1 ed; 2017.
- [47] Freedman GE. srvyr: 'dplyr'-like syntax for summary statistics of survey data. R package version 0.3.4 ed; 2019.
- [48] Lumley T. Survey: analysis of complex survey samples R package version 3.35-1 ed; 2019.
- [49] Long J. _jtools: analysis and presentation of social scientific data_. R package version 2.0.1 ed; 2019.
- [50] Wei T, Simko V. R package “corrplot”: visualization of a correlation matrix (Version 0.84). Version 0.84 ed; 2017.
- [51] Fox J, Weisberg S. *An R companion to applied regression*. 2nd ed. Thousand Oaks CA: Sage; 2011.
- [52] Mead P, Hinckley A, Hook S, Beard CB. TickNET—A collaborative public health approach to tickborne disease surveillance and research. *Emerg Infect Dis* 2015;21(9):1574–7.
- [53] Barone SR, Bohrer SS, Erhardt WA. Parental knowledge of and attitudes toward LYMERix (recombinant Osp-a Lyme vaccine). *Clin Pediatr (Phila)* 2002;41(1):33–6.
- [54] Nolan K, Mauer MP. An evaluation of a Lyme disease prevention program in a working population. *Am J Health Promot* 2006;20(6):379–82.
- [55] Nawrocki CC, Hinckley AF. Experiences with tick exposure, Lyme disease, and use of personal prevention methods for tick bites among members of the U.S. population, 2013–2015. *Ticks Tick-borne Dis* 2021;12(1).
- [56] Smith PJ, Chu SY, Barker LE. Children who have received no vaccines: who are they and where do they live? *Pediatrics* 2004;114(1):187–95.
- [57] Williams WW, Lu P-J, O'Halloran A, Kim DK, Grohskopf LA, Pilishvili T, et al. Surveillance of vaccination coverage among adult populations – United States, 2015. *MMWR Surveill Summ* 2017;66(11):1–28.
- [58] Williams WW, Lu PJ, O'Halloran A, Bridges CB, Kim DK, Pilishvili T, et al. Vaccination coverage among adults, excluding influenza vaccination – United States, 2013. *MMWR Morb Mortal Wkly Rep* 2015;64:95–102.
- [59] Smith PJ, Humiston SG, Parnell T, Vannice KS, Salmon DA. The association between intentional delay of vaccine administration and timely childhood vaccination coverage. *Public Health Rep* 2010;125(4):534–41.
- [60] Salmon DA, Moulton LH, Omer SB, deHart MP, Stokley S, Halsey NA. Factors associated with refusal of childhood vaccines among parents of school-aged children: a case-control study. *Arch Pediatr Adolesc Med* 2005;159(5):470. <https://doi.org/10.1001/archpedi.159.5.470>.
- [61] Jacobson RM, St. Sauver JL, Griffin JM, MacLaughlin KL, Finney Rutten LJ. How health care providers should address vaccine hesitancy in the clinical setting: evidence for presumptive language in making a strong recommendation. *Hum Vaccin Immunother* 2020;16(9):2131–5.
- [62] Shaffer L. Inner workings: Lyme disease vaccines face familiar challenges, both societal and scientific. *Proc Natl Acad Sci U S A* 2019;116(39):19214–7.
- [63] Lu P-J, Yankey D, Fredua B, O'Halloran AC, Williams C, Markowitz LE, et al. Association of provider recommendation and human papillomavirus vaccination initiation among male adolescents aged 13–17 Years—United States. *J Pediatr* 2019;206:33–41.e1.
- [64] Gargano LM, Herbert NL, Painter JE, Sales JM, Morfaw C, Rask K, et al. Impact of a physician recommendation and parental immunization attitudes on receipt or intention to receive adolescent vaccines. *Hum Vaccin Immunother* 2013;9(12):2627–33.
- [65] Salmon DA, Dudley MZ, Glanz JM, Omer SB. Vaccine hesitancy: causes, consequences, and a call to action. *Vaccine* 2015;33(Suppl 4):D66–71.
- [66] Liao Q, Cowling BJ, Lam WW, Fielding R. Factors affecting intention to receive and self-reported receipt of 2009 pandemic (H1N1) vaccine in Hong Kong: a longitudinal study. *PLoS One* 2011;6(3).
- [67] Gidengil CA, Parker AM, Zikmund-Fisher BJ. Trends in risk perceptions and vaccination intentions: a longitudinal study of the first year of the H1N1 pandemic. *Am J Public Health* 2012;102(4):672–9.
- [68] Maurer J. Inspecting the mechanism: a longitudinal analysis of socioeconomic status differences in perceived influenza risks, vaccination intentions, and vaccination behaviors during the 2009–2010 influenza pandemic. *Med Decis Making* 2016;36(7):887–99.
- [69] Devchand Roshni, Koehler Laura, Hook Sarah, Marx Grace, Hooks Holley, Schwartz Amy, et al. Understanding consumer and clinician perceptions of a potential Lyme disease vaccine. *Health Education Research* 2021.

Question	Response options
If a Lyme disease vaccine were available, would you get vaccinated?	Yes No Don't know/not sure
How concerned are you about the safety of a Lyme disease vaccine?	Not at all concerned Somewhat concerned Very concerned Don't know/not sure
How much would the cost of a Lyme disease vaccine affect your decision to get vaccinated?	Not at all Some A lot Don't know/not sure
How much would a positive recommendation from your doctor affect your decision to get vaccinated?	Not at all Some A lot Don't know/not sure
Has anyone in your household ever been diagnosed with LD by a health care professional?	Yes No Not sure
How concerned are you about getting LD in the future?	Not at all concerned Somewhat concerned Very concerned Don't know/not sure
In the months April-October, do you spend time in or near places where ticks could get on you (for example, wooded or brushy areas, whether in your yard, other yards, or recreational areas)?	Yes, daily Yes, weekly Yes, monthly Yes, less than once a month No Don't know/not sure
Which of the following measures do you take to prevent ticks from getting on you? (Check all that apply)	Apply insect repellent Check for ticks Use special clothing Use sprays in your yard Other measures None of these
How confident are you that these measures can prevent LD?	Very confident Somewhat confident Not at all confident Don't know/not sure
Where do you most often get information about Lyme disease? (choose one)	Doctor, nurse, or other medical professional Naturopath or chiropractor Friends or family members Google or other internet search engines Health websites Social media sites Other

Appendix: Evaluating public acceptability of a potential Lyme disease vaccine in the United States

How confident are you that recommended vaccines benefit people?	Very confident Somewhat confident Not at all confident Don't know/not sure
Where do you usually get vaccines? (choose one)	Doctor's office, clinic, or hospital Pharmacy or drug store Health department Workplace School clinic Other Don't know I do not get vaccines
<p><i>* Questions for the parent survey were identical, but phrased with the child as vaccinee (e.g., "If a LD vaccine were available, would you vaccinate your child?")</i></p>	

3 **Table A2. Confounders identified *a priori* and adjusted for in multinomial logistic regression models**

Model	Confounders											
	Vaccinee age category	Gender	State	Race	Education	Metro status	Healthcare provider recommendation	Past LD diagnosis in household	Concern about future LD diagnosis	Time spent in tick habitat	Current use of LD prevention measures	General confidence in vaccines
Vaccinee age category*												
Gender*												
State	X				X							
Race						X						
Education	X	X	X	X		X						
LD vaccine safety concerns	X	X			X		X	X	X	X	X	X
HCP influence on LD vaccination	X	X			X			X	X	X	X	X
LD vaccine cost concerns	X	X			X			X	X	X		X

4 *Unadjusted models

5

6 **Table A3. Observed and weighted respondent characteristics, N = 3206**

Characteristic	N	Unweighted %	Weighted % (95% CI) *
Demographics			
Gender**			
Female	1878	59	54
Male	1328	41	46
Age category** (years)			
< 18	246	8	15
18-44	772	24	33
45-64	1225	38	34
65+	963	30	17
State			
Connecticut	679	21	20
Maryland	808	25	27
Minnesota	998	31	20
New York	721	23	33
Race			
White	2852	90	85 (84, 86)
Non-white	322	10	15 (14, 16)
Education			
Some college or less	1248	39	35 (33, 36)
Bachelor's degree or higher	1941	61	65 (64, 67)
Metropolitan status			
Large central metropolitan area	674	21	28
Other	2532	79	72
LD history, attitudes, and practices			
Past LD diagnosis in household			
Yes	640	20	18 (17, 19)
No	2563	80	82 (81, 83)
Concern about future LD diagnosis			
Yes	2813	88	86 (85, 86)
No	391	12	14 (14, 15)
Spend time in tick habitat			
At least weekly	2376	74	71 (70, 73)
Monthly or less	828	26	29 (27, 30)
Currently use LD prevention measures			
Yes	2948	92	92 (91, 93)
No	258	8	8 (7, 9)
Confidence in LD prevention measures			
Yes	2041	70	70 (68, 71)
No	896	30	30 (29, 32)

Appendix: Evaluating public acceptability of a potential Lyme disease vaccine in the United States

Confidence in general vaccines

Yes	3022	94	94 (93, 95)
No	182	6	6 (5, 7)

LD vaccine attitudes

Willing to receive LD vaccine

Yes	2098	65	64 (62, 65)
No	190	6	7 (6, 8)
Don't know	918	29	30 (28, 31)

LD vaccine safety concerns

Yes	2257	70	71 (70, 72)
No	948	30	29 (28, 30)

HCP influence on LD vaccination

Yes	2858	89	89 (88, 89)
No	348	11	11 (11, 12)

LD vaccine cost concerns

Yes	2036	64	63 (62, 65)
No	1168	36	37 (35, 38)

7 * County distributions of gender and age were used for post-stratification; as such, these point estimates are fixed at the
8 population values and have no associated interval estimate. Because state and metropolitan status are based on county
9 population totals, these point estimates are also fixed.

10 **Gender and age categories represent the potential vaccinee, i.e., adult respondents and the children for whom parents
11 responded.

12

13

14 **Section 1. Heckman-type selection models**

15 We evaluated non-random missingness in our outcome variable, willingness to receive a LD vaccine, in
16 relation to non-response (i.e., selection bias) using Heckman-type selection models, also called
17 generalized Tobit models [39-41]. Heckman-type selection models correct for selection bias when
18 nonparticipation is determined both by observed and by unobserved factors. Performance depends on
19 the availability of selection variables that determine survey participation but do not independently
20 affect the outcome of interest. Heckman models use two steps to first model the selection process using
21 one or more independent selection variables and then model the outcome equation (i.e., the regression
22 equation for the outcome of interest). The key feature of Heckman-type selection models is that a
23 correlation between the unobserved error terms in the selection equation and outcome equation is
24 estimated (ρ). The coefficient of the inverse Mill's ratio represents the covariance between the error
25 terms, and has an associated p-value. These results of the two-step process indicate whether selection
26 bias is present and, if so, a correction factor incorporating the coefficient of the inverse Mill's ratio is
27 applied to results.

28 We chose two selection variables, presence of children in the household and household member count,
29 under the assumption that these variables were predictive of participation in the survey, but unrelated
30 to the outcome, willingness to receive a LD vaccine. For example, those with children in the household
31 and/or higher numbers of household members may not have time to participate in a voluntary survey.
32 All variables in the selection equation must be available for all sampled individuals, regardless of
33 participation. Independent variables for all sampled households were purchased from the marketing
34 firm from which addresses were purchased. The selection equation included the following:

35 **Selection ~ endemicity + property type + household income + presence of children in**
36 **household + household member count**

37 The outcome equation includes the independent variables from the selection equation, excluding the
38 selection variables. The outcome equation included the following:

39 **Vax decision ~ endemicity + property type + household income**

40 Table A4 shows the results of the Heckman-type selection models using the two-step process. Of note, ρ
41 = 0.7 and the coefficient of the inverse Mill's ratio is 0.4305 ($p = 0.5307$), meaning that the data are
42 consistent with no selection bias (i.e., the null hypothesis that the errors are uncorrelated cannot be
43 rejected).

44 These results are limited by the fact that only variables available for the entire sample could be used in
45 the evaluation. Further, the accuracy of these variables typically used for marketing research are
46 questionable, plus some records were missing observations for these variables. Lastly, our assumption
47 that the selection variables, presence of children in household and household member count, are
48 unrelated to the willingness to be vaccinated outcome is somewhat tenuous. For example, because
49 children are one of the groups at highest risk for LD, parents with children in the household may be
50 more likely to participate in a survey about a LD vaccine and also express willingness to vaccinate their
51 children.

52

53

Table A4. Results of Heckman-type selection models

<i>Dependent variable:</i> Willingness to receive LD vaccine (0= No/DK; 1 = Yes)	
Terms	Coefficient (95% CI)
Endemicity:	-0.0223
Non-endemic	(-0.1285, 0.0840)
Property type:	0.0690
Single family dwelling unit	(-0.0825, 0.2206)
Household income:	0.0709
> \$70K	(-0.0868, 0.2286)
Constant	-0.2494 (-2.4066, 1.9078)
Observations	34,667
R ²	0.0006
Adjusted R ²	-0.0009
Log Likelihood	
Akaike Inf. Crit.	
rho	0.6958
Inverse Mills Ratio	0.4305 (0.5307)


54

55 ¹The selection equation for the Heckman selection model used presence of children and household member count
 56 as selection variables, i.e., instrumental variables

57 Note: *p **p ***p<0.01

58

59 **Figure A1. Respondents' primary source for LD information, by potential LD vaccination decision,**
60 **weighted % and 95% confidence interval***




61

62 *95% confidence interval shown in the black bars

63

64 **Figure A2. Respondents' primary location for receiving vaccination, by potential LD vaccination**
 65 **decision, weighted % and 95% confidence interval***



66

67 *95% confidence interval shown in the black bars